

A local Fourier analysis for additive Schwarz smoothers

Álvaro Pé de la Riva^a, Carmen Rodrigo^{a,*}, Francisco J. Gaspar^a, James H. Adler^b, Xiaozhe Hu^b,
Ludmil Zikatanov^c

^a IUMA and Applied Mathematics Department, University of Zaragoza, Spain

^b Tufts University, Medford, MA, USA

^c Penn State, University Park, PA, USA

ARTICLE INFO

Keywords:

Local Fourier analysis
Multigrid methods
Additive Schwarz smoothers
High-order finite element methods
Saddle point problems

ABSTRACT

In this work, a local Fourier analysis is presented to study the convergence of multigrid methods based on additive Schwarz smoothers. This analysis is presented as a general framework which allows us to study these smoothers for any type of discretization and problem. The presented framework is crucial in practice since it allows one to know a priori the answer to questions such as what is the size of the patch to use within these relaxations, the size of the overlap, or even the optimal values for the weights involved in the smoother. Results are shown for a class of additive and restricted additive Schwarz relaxations used within a multigrid framework applied to high-order finite-element discretizations and saddle point problems, which are two of the contexts in which these types of relaxations are widely used.

1. Introduction

Multigrid methods are among the best-known iterative solution techniques due to their demonstrated high efficiency for a wide range of problems. They accelerate the convergence of classical iterative methods by combining them with a coarse-grid correction technique. The design of efficient multigrid methods, however, depends crucially on the choice of their components. One of the most important ingredients of a multigrid algorithm is the so-called smoother or relaxation procedure, which often consists of a classical iterative method such as Jacobi or Gauss-Seidel.

Within a multigrid framework, a natural extension of point-wise smoothers is patch-wise smoothers. In order to apply such a relaxation, the computational domain is divided into small (overlapping or non-overlapping) patches, and then, one smoothing step consists of solving local problems on each patch one-by-one either in a Jacobi-type or Gauss-Seidel-type manner. This results in an additive or multiplicative Schwarz smoother, respectively. One of the best-known multigrid methods based on this type of relaxation was proposed by Vanka in [1] for solving the steady-state incompressible Navier-Stokes equations in primitive variables, discretized by a finite-volume scheme on a staggered grid. The computational domain is divided into cells with pressure nodes at the cell centers and velocity nodes at the cell faces. The smoothing procedure is a so-called symmetric coupled Gauss-Seidel

technique (SCGS), which consists of solving local problems for each cell involving all the unknowns located at the cell. This is done cell by cell in a Gauss-Seidel-type manner and, therefore, can be viewed as a multiplicative Schwarz iteration.

Additive Schwarz-type iteration methods have been studied in [2] as smoothers in a multigrid method for saddle point problems. It is shown that, under suitable conditions, the iteration can be interpreted as a symmetric inexact Uzawa method. Restrictive additive Schwarz methods (RAS) were introduced in [3] as an efficient alternative to the classical additive Schwarz preconditioners. They introduced a simple change to the additive scheme by removing the overlap in the interpolation operator, becoming more attractive as they reduce the communication time between processors and usually the overall computation time. Convergence of RAS methods was proven in [4], where it was shown that this method reduces communication time while maintaining the most desirable properties of the classical Schwarz methods. RAS preconditioners are widely used in practice and are implemented in several software packages. In [5], a restrictive Schwarz method was proposed as a smoother for solving the Stokes equations. It was observed that this smoother achieves comparable convergence rates to the multiplicative version, while being computationally less expensive per iteration. In general, the class of additive and restricted additive Schwarz smoothers is characterized by their ability to deal with high-

* Corresponding author.

E-mail addresses: apedelariva@unizar.es (Á. Pé de la Riva), carmenr@unizar.es (C. Rodrigo), fjgaspar@unizar.es (F.J. Gaspar), james.adler@tufts.edu (J.H. Adler), xiaozhe.hu@tufts.edu (X. Hu), ludmil@psu.edu (L. Zikatanov).

<https://doi.org/10.1016/j.camwa.2023.12.039>

Received 14 July 2023; Received in revised form 23 December 2023; Accepted 31 December 2023

order discretizations, saddle point problems, and equations where the terms grad-div or curl-curl dominate. Thus, in this work, we aim to study such relaxations within a multigrid framework.

Local Fourier analysis (LFA), or local mode analysis, is a commonly used approach for analyzing the convergence properties of geometric multigrid methods. In this analysis an infinite regular grid is considered and boundary conditions are not taken into account. LFA was introduced by Brandt in [6], and afterwards extended in [7]. A good introduction can be found in the paper by Stüben and Trottenberg [8] and in the books by Wesseling [9], Trottenberg et al. [10], and Wienands and Joppich [11]. It is the main quantitative analysis for the convergence of multilevel algorithms, and results in a very useful tool for the design of multigrid methods. Moreover, in [12] it has been recently proved that under standard assumptions LFA is a rigorous analysis, providing the exact asymptotic convergence factors of the multigrid method.

LFA for multiplicative Schwarz smoothers was first performed in [13] for the staggered finite-difference discretization of the Stokes equations, and in [14] for a mixed finite-element discretization of the Laplace equation. In [15], an LFA for multiplicative Schwarz smoothers on triangular grids is presented, and in [16], the analysis for such overlap block smoothers is performed on rectangular grids for finite-element discretizations of the grad-div, curl-curl and Stokes equations.

Whereas LFA has been widely applied to multiplicative Schwarz relaxations, few works on additive Schwarz smoothers can be found in the literature. Whereas the Fourier modes are eigenfunctions of the multiplicative Schwarz smoothers (see [16] for a rigorous proof), this statement is not true for additive Schwarz smoothers. In [17], an LFA to analyze a multicolored version of an additive Schwarz smoother for a curl-curl model problem was proposed. A non-standard LFA to analyze this type of smoothers for the Stokes equations with P2-P1 and Q2-Q1 discretizations is considered in [18]. This analysis was also used in [19] in order to study an additive Vanka-type smoother within a multigrid framework for the Poisson equation discretized by a standard finite-difference scheme. The analysis developed in both references can be seen as a particular case of the general framework analysis presented in this work, which allows us to study this class of smoothers for any type of discretization and problem.

Finally, we note that a non-standard LFA technique to predict the convergence rate of multigrid solvers for problems involving random and jumping coefficients was proposed in [20]. Similar analysis has been recently introduced independently in [21–23]. The novelty of this new approach lies in the use of a specific basis of the Fourier space, rather than the standard basis which is based on the Fourier modes. This is the approach that we consider in this work and, as it will be shown, it allows us to propose an LFA to study the convergence of multigrid methods based on additive Schwarz smoothers for any discretization and problem. This general framework is crucial in practice, since it can answer questions such as what the appropriate size of the patch to use is, how to determine the size of the overlap, what the optimal values for the weights involved in the smoother are, and whether or not the restricted additive version is the best choice.

The rest of the paper is organized as follows. In Section 2, the additive and restricted additive Schwarz smoothers are introduced together with their corresponding algorithms. Section 3 is devoted to presenting the basis for the LFA performed in this work. Sections 4 and 5 deal with the application of the proposed local Fourier analysis to high-order finite-element discretizations for scalar problems and saddle point type problems, respectively. Finally, a summary and concluding remarks are given in Section 6.

2. Additive Schwarz methods

In this section we introduce the additive Schwarz methods used to solve a linear system of n algebraic equations, $Au = b$. Consider a decomposition of the unknowns into subsets B_i , $i = 1, \dots, s$, such that each unknown in vector u is included in at least one block B_i . For each sub-

set of unknowns B_i , let $V_i : R^n \rightarrow R^{n_i}$ be the mapping from the global vector of unknowns to the ones given in B_i , where n_i is the size of B_i . Here, V_i is represented by a rectangular $n_i \times n$ matrix and, with a slight abuse of notation, we also refer to this matrix as V_i . Its transpose, V_i^T , is a prolongation operator from R^{n_i} to R^n by padding it with zeros. Then, $A_i = V_i A V_i^T$ is the local matrix corresponding to the unknowns in B_i , that is, the restriction of A to B_i . For each block B_i , the following small local system has to be solved at the j th iteration,

$$A_i \delta u_i^j = V_i(b - Au^j),$$

where δu_i^j denotes the vector of corrections for the unknowns involved in block B_i .

These corrections can be computed via a multiplicative or an additive Schwarz method. Although multiplicative Schwarz methods provide better asymptotic convergence factors, in general their additive counterparts have lower cost per iteration since they are more parallelizable. This is due to the fact that the corrections provided by solving the other local systems are not required to compute the corrections associated with a given block. The algorithm of the additive Schwarz method is given as follows:

Algorithm 1: Additive Schwarz method.

Input: u^j . **Output:** u^{j+1} .
1 $r = b - Au^j$
2 **for** $i = 1 : s$ **do**
3 **Solve:** $A_i \delta u_i^j = V_i r$.
4 $u^{j+1} = u^j + \sum_{i=1}^s V_i^T D_i \delta u_i^j$,

In Algorithm 1, D_i is, in the general case, a diagonal weighting matrix. The diagonal entries of this matrix are usually taken as the inverse of the number of blocks sharing the corresponding unknown, but can be tuned to improve the convergence factor of the method.

The restricted additive Schwarz method is based on the use of prolongation operators, \tilde{V}_i^T , in such a way that each entry of the vector unknown u occurs in $\tilde{V}_i^T u$ for exactly one i . Therefore, \tilde{V}_i^T is a matrix of zeros and ones of the same size as V_i^T , such that the correction to each unknown is obtained from a single block. This means that prolongation operators \tilde{V}_i^T are introduced in such a way that

$$\sum_{i=1}^s \tilde{V}_i^T V_i = I,$$

is satisfied. Then, the algorithm of the restricted additive Schwarz method is:

Algorithm 2: Restricted additive Schwarz method.

Input: u^j . **Output:** u^{j+1} .
1 $r = b - Au^j$
2 **for** $i = 1 : s$ **do**
3 **Solve:** $A_i \delta u_i^j = V_i r$.
4 $u^{j+1} = u^j + \sum_{i=1}^s \tilde{V}_i^T D_i \delta u_i^j$,

When using these types of smoothers for solving a particular problem, some key questions arise. First, what should the size of the blocks be? Second, how large should the overlap be? Finally, the weights involved in the method are input parameters, and are extremely important to obtain an efficient and robust solver. Thus, determining the optimal values is a big concern. In the next section, we present a general framework based on LFA that answers these questions, allowing us to know a priori the type of additive Schwarz method to use.

3. Local Fourier analysis

In this section, we describe LFA in a setting which allows us to estimate the multigrid convergence factors by using additive Schwarz

methods as smoothers. Standard LFA assumes that Fourier modes are eigenfunctions of some of the operators involved in the multigrid algorithm, such as the discrete operator and the relaxation procedure. However, we mention three examples where this conventional analysis is not directly applicable. The first exception consists of problems with non-constant coefficients, as models involving random and jumping coefficients. Secondly, discretizations with different stencils at distinct grid-points, appearing for example in high-order finite-element methods also pose challenges to the classical approach. Finally, standard LFA is not applicable when Fourier modes are not eigenfunctions of the smoother considered in the multigrid algorithm. Additive Schwarz methods fall in this latter category.

In [20], a non-standard LFA technique to predict the convergence rate of multigrid solvers for problems involving random and jumping coefficients was proposed. While the analysis in that work was applied only to problems with non-constant coefficients, it can be also applied to high-order finite-element discretizations. In fact, the special LFA performed in [24] to study multigrid methods for quadratic finite-element methods can be seen as a particular case of the analysis presented in [20]. In addition, as will be shown here, it can also be applied for analyzing multigrid methods based on additive Schwarz smoothers. Next, we briefly describe this analysis, and refer to [20] for more details.

Given an infinite and regular grid, \mathcal{G}_h with grid size h , Fourier modes are defined by $\varphi_h(x, \theta) = e^{i\theta x}$, with $x \in \mathcal{G}_h$ and $\theta \in \Theta_h := (\pi/h, \pi/h]$. The Fourier modes span the so-called Fourier space:

$$\mathcal{F}_h(\mathcal{G}_h) = \text{span}\{\varphi_h(x, \theta) \mid \theta \in \Theta_h\}. \quad (1)$$

The main idea in the analysis presented in [20] is to consider a specific basis of the Fourier space, rather than the standard basis which is based on the Fourier modes. For this purpose, we consider a splitting of \mathcal{G}_h into m^d infinite subgrids, where d is the dimension of the problem. We then take a fixed window of size $\underbrace{m \times \dots \times m}_d$ and its periodic extension. The size of the window has to be chosen appropriately, which we do in the context of additive Schwarz smoothers. For the sake of simplicity, we consider the same size m in each direction. We note that the analysis still holds even if they are different. Once m is fixed, for every $\mathbf{k} = (k_1, \dots, k_d)$, $k_1, \dots, k_d = 0, \dots, m-1$, subgrid, $\mathcal{G}_h^{\mathbf{k}}$, is defined as follows:

$$\mathcal{G}_h^{\mathbf{k}} = \{\mathbf{k}h + (l_1, \dots, l_d)mh \mid l_1, \dots, l_d \in \mathbb{Z}\}. \quad (2)$$

For each frequency $\theta^0 \in \Theta_{mh} := (-\pi/mh, \pi/mh]^d$, we introduce the following grid functions:

$$\psi_h^{\mathbf{k}}(\theta^0, \mathbf{x}) = \varphi_h(\theta^0, \mathbf{x}) \chi_{\mathcal{G}_h^{\mathbf{k}}}. \quad (3)$$

As in [20], the Fourier space spanned by these functions,

$$\mathcal{F}_h^{m^d}(\theta^0) = \text{span}\{\psi_h^{\mathbf{k}}(\theta^0, \cdot), \mathbf{k} = (k_1, \dots, k_d), k_1, \dots, k_d = 0, \dots, m-1\}, \quad (4)$$

is the same Fourier space generated by the standard Fourier modes,

$$\text{span}\{\varphi_h(\theta_k^0, \cdot), \theta_k^0 = \theta^0 + \mathbf{k} \frac{2\pi}{mh}, \mathbf{k} = (k_1, \dots, k_d), k_1, \dots, k_d = 0, \dots, m-1\}. \quad (5)$$

Due to the relation between the grid-functions of the given new Fourier basis and the standard Fourier modes, it can be shown that, taking a big enough window size, the two-grid operator,

$$M_h = (I_h - P_{2h}^h A_{2h}^{-1} R_h^{2h} A_h) S_h^\nu, \quad (6)$$

satisfies the following invariance property for any frequency $\theta^0 \in \Theta_{mh}$,

$$M_h : \mathcal{F}_h^{m^d}(\theta^0) \rightarrow \mathcal{F}_h^{m^d}(\theta^0).$$

In (6), P_{2h}^h and R_h^{2h} are the prolongation and restriction operators, A_h and A_{2h} are the fine- and coarse-grid operators, I_h is the identity, S_h is the smoothing operator, and ν is the number of smoothing steps.

When the usual analysis can be applied, and a standard coarsening $H = 2h$ is considered, the Fourier space has dimension 2^d , yielding the so called $2h$ -harmonics spaces. In this case, $m = 2$ in (4), the size of the window is 2^d , and both analyses coincide. The advantage of using the grid-functions in (3) is that it allows one to study problems for which the analysis based on the Fourier modes is not applicable. For example, using $m = 2$ one could study the multigrid convergence for quadratic finite-element methods. As we will see in the next section, one can study finite-element methods of arbitrary order just taking a bigger value of m . Additionally, we will show that Additive Schwarz methods can be analyzed by considering the grid-functions in (3), and choosing an adequate window size, i.e., an adequate value of m .

4. High-order finite-element methods for scalar problems

As a model problem, we consider the Poisson equation in a d -dimensional domain, $\Omega = (0, 1)^d$, with homogeneous Dirichlet boundary conditions,

$$\begin{aligned} -\Delta u &= f, \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega. \end{aligned} \quad (7)$$

The variational formulation of this problem reads as follows: Find $u \in H_0^1(\Omega)$ such that

$$a(u, v) = (f, v), \quad \forall v \in H_0^1(\Omega),$$

where

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v \, dx, \quad \text{and} \quad (f, v) = \int_{\Omega} f v \, dx.$$

Let \mathcal{T}_h be a partition of the domain $\Omega \subset \mathbb{R}^d$, and associate to \mathcal{T}_h the high-order finite-element space, $V_{hp} \subset H_0^1(\Omega)$ defined as $V_{hp} = \{u_h \in H_0^1(\Omega) \mid u_h|_T \in \mathbb{Q}_p, \forall T \in \mathcal{T}_h\}$, where \mathbb{Q}_p is the space of polynomials up to total degree p on each variable. Thus, the Galerkin approximation of the variational problem is given by: Find $u_h \in V_{hp}$, such that

$$a(u_h, v_h) = (f, v_h), \quad \forall v_h \in V_{hp}. \quad (8)$$

The solution of the Galerkin approximation problem is a linear combination, $u_h = \sum_{i=1}^{n_h} u_i \varphi_i$, where $\{\varphi_1, \dots, \varphi_{n_h}\}$ is a basis for V_{hp} and $\dim V_{hp} = n_h$. In order to compute $\mathbf{u} = (u_1, \dots, u_{n_h})^T$, the following linear system must be solved: $\mathbf{A}\mathbf{u} = \mathbf{b}$, where $A = (a_{ij}) = (a(\varphi_j, \varphi_i))_{i,j=1}^{n_h}$ is the so-called stiffness matrix and $\mathbf{b} = (b_i) = (f, \varphi_i)_{i=1}^{n_h}$ is the right-hand side vector.

The goal now is to demonstrate the utility of the analysis presented in the previous section for the design of efficient geometric multigrid methods based on additive Schwarz relaxation schemes. For this purpose, we consider an h -multigrid, i.e., we keep the polynomial order unchanged in the geometric mesh hierarchy, combined with the canonical high-order restriction and prolongation operators. There are, however, other possibilities to build a multilevel hierarchy for high-order discretized problems. Another option is to use a p -multigrid technique, which consists of constructing the coarse problems within the multigrid algorithm by reducing the polynomial degree of the finite-element space. This approach can be combined with a standard h -multigrid on the coarsest level $p = 1$. Recently in [25], a two-level method based on multiplicative Schwarz methods was proposed such that a discretization of arbitrary order p is considered in the first level whereas the second level consists of a linear discretization ($p = 1$) of the problem. Although we only present results here for the h -multigrid case, the analysis proposed in this work can be applied to any other alternative for high-order discretizations, such as those approaches mentioned above.

Table 1

Linear finite-element discretization, $p = 1$, for the 1D Poisson problem. Two-grid asymptotic convergence factors applying $\nu = 1$ smoothing steps of an additive Schwarz method. Block-size, (k) , ranges from 2 to 7 and the size of the overlap, ov , from 1 to $k - 1$. Natural weights are taken in the smoother.

Overlap (ov)	Block-size (k)					
	2	3	4	5	6	7
1	0.33	0.99	0.40	0.50	0.43	0.50
2	-	0.33	0.20	0.50	0.29	0.25
3	-	-	0.20	0.50	0.21	0.99
4	-	-	-	0.20	0.14	0.25
5	-	-	-	-	0.14	0.33
6	-	-	-	-	-	0.14

4.1. One-dimensional case

First, we consider the simplest case when $d = 1$, the computational domain is the interval, $\Omega = (0, 1)$, and the corresponding two-point boundary value problem is $-u''(x) = f(x)$, $x \in \Omega$, $u(0) = u(1) = 0$. We consider linear finite-element methods, i.e., $p = 1$, and perform LFA based on infinite subgrids to estimate the convergence factor of the multigrid method using additive Schwarz methods as the relaxation. In additive Schwarz, we vary the size of the block, k , from 2 to 7 and the size of the overlap, ov , from $ov = 1$ to the maximum overlap $ov = k - 1$. The diagonal elements of the weighting matrix, D_i , in Algorithm 1 are taken to be the natural weights, i.e., the reciprocal of the number of blocks in which each unknown appears. For example, if the size of the block is $k = 3$ and the size of the overlap is $ov = 1$, the weights for the three unknowns of a block are $1/2, 1, 1/2$. In Table 1, we show the two-grid asymptotic convergence factors provided by LFA with only one smoothing iteration, $\nu = 1$, for different block-sizes and overlap sizes, considering a linear finite-element discretization of the 1D Poisson equation. We do not show the experimentally computed asymptotic convergence factors because they are exactly the same as those provided by LFA.

From the results presented in Table 1, we observe that a bigger block-size or overlap among blocks does not always provide a better asymptotic convergence factor.

Next, we consider the restrictive additive Schwarz method as the smoother within the multigrid framework. Now, the natural weights will be one or zero. For example, if the size of the block is $k = 3$ and the size of the overlap is $ov = 1$, the weights for the three unknowns of the block are 1, 1, 0. In Table 2, we show the asymptotic convergence factors provided by our two-grid analysis by using one smoothing step of the restrictive additive Schwarz method with natural weights. Block-sizes ranging from 2 to 7 and all the possible overlap among the blocks are considered. Again, the experimentally computed asymptotic convergence factors are the same as those provided by the LFA, and thus not included in the table.

The obtained results are not good in general, observing even no convergence of the multigrid method for some combinations of block-size and size of the overlap. The advantage of having the LFA tool provided in this work, however, is that this analysis can find the optimal weights to improve the performance of the multigrid method. Hence, in Table 3, we show the asymptotic convergence factors together with the optimal weights provided by the analysis. Again, the two-grid analysis is based on only one smoothing step of the restrictive additive Schwarz methods with optimal weights, and different block-sizes and sizes of the overlap among the blocks are considered.

Comparing the results in Tables 2 and 3, we observe a significant improvement of the asymptotic convergence factors when appropriate non-trivial weights are used. Once again, better asymptotic convergence factors are not directly related to bigger block-sizes or larger overlap among blocks.

Table 2

Linear finite-element discretization, $p = 1$, for the 1D Poisson problem. Two-grid asymptotic convergence factors applying $\nu = 1$ smoothing steps of a restrictive additive Schwarz method with natural weights. Block-size, (k) , ranges from 2 to 7 and the size of the overlap, (ov) , from 1 to $k - 1$.

Overlap	Block-size					
	2	3	4	5	6	7
1	0.75	1.00	0.40	0.50	0.43	0.50
2	-	1.00	0.60	0.50	0.57	0.44
3	-	-	0.87	0.87	0.28	1.00
4	-	-	-	1.00	0.71	0.44
5	-	-	-	-	0.92	1.00
6	-	-	-	-	-	1.00

LFA codes corresponding to previous tables are available open-source [26].

The proposed LFA also lets us study the sensitivity of the convergence factors of the multigrid method with respect to small changes in the weights. In Fig. 1, we display the two-grid convergence factor versus the weight chosen for two combinations of block and overlap size: $k = 4$, $ov = 2$ and $k = 5$, $ov = 3$. In both cases, when weights are close to optimal, the difference in their convergence factors is insignificant. Therefore, selecting any weight close to optimal will be equally effective.

Our analysis, though, allows us to extend LFA to high-order discretizations. Here, we fix the size of the block and the overlap and we vary the polynomial order in the finite-element space. An element-based additive Schwarz method is considered, where each block contains all the basis functions with support in the corresponding element. Therefore, the size of the blocks is chosen as $p + 1$, where p denotes the polynomial degree. The size of the overlap is fixed to be the minimum, i.e., $ov = 1$. In Table 4, we show the asymptotic convergence factors estimated by the two-grid analysis by using one smoothing step of the additive Schwarz method (AS) and one step of the restricted additive Schwarz method (RAS). The polynomial degrees range from $p = 2$ to $p = 8$.

For each polynomial degree, p , we find that the restricted additive Schwarz smoother achieves comparable convergence rates to the additive Schwarz smoother, having better properties in terms of scalability and applicability to high-performance computing. We conclude that the restricted additive Schwarz method is a favorable smoother for high-order discretizations of elliptic equations. The authors are not aware of any work comparing these approaches.

Remark 1. The size of the window, m , in the Fourier analysis is the least common multiple of $2p$ and the minimum number $(k - ov)j$ such that $(k - ov)j > k$, with j being any natural number. In the particular case that the size of the block $k = p + 1$ and $ov = 1$, the size of the block is the least common multiple of $2p$ and the minimum number pj such that $pj > p + 1$. For example, if $p = 2$ the size of the windows is $m = 4$.

4.2. Two-dimensional case

We now consider the two-dimensional case of model problem (7). We follow the same element-based smoothing strategy using additive Schwarz methods. Thus, the size of local linear systems will be $k = (p + 1)^2$ and the overlap among the blocks in both directions is still one, $ov = 1$. In Table 5, we show the asymptotic convergence factors predicted by the two-grid analysis by using $\nu = 2$ smoothing steps of the corresponding AS and the RAS methods with natural weights. The polynomial degree ranges from $p = 1$ to $p = 8$.

Comparing the results of both smoothers in Table 5, we conclude that the restrictive case is better from a computational cost point of

Table 3

Linear finite-element discretization, $p = 1$, for the 1D Poisson problem. Two-grid asymptotic convergence factors (and optimal weights in parenthesis) applying one iteration of the restrictive additive Schwarz method with optimal weights. Block-size, (k), ranges from 2 to 7 and the size of the overlap, (ov), from 1 to $k - 1$.

Ov	Block-size					
	2	3	4	5	6	7
1	0.45 (0.6)	0.34 (0.67)	0.17 (0.83)	0.20 (0.8)	0.18 (0.82)	0.20 (0.8)
2	-	0.37 (0.68)	0.15 (0.71)	0.20 (0.8)	0.22 (0.78)	0.18 (0.82)
3	-	-	0.43 (0.71)	0.34 (0.66)	0.16 (0.84)	0.34 (0.67)
4	-	-	-	0.36 (0.66)	0.20 (0.7)	0.18 (0.82)
5	-	-	-	-	0.40 (0.7)	0.34 (0.66)
6	-	-	-	-	-	0.34 (0.66)




Fig. 1. Two grid convergence factors predicted by LFA applying one smoothing step of the restrictive additive scheme for different weights (a) Block size $k = 4$ and size of the overlap $ov = 2$ (b) Block size $k = 5$ and size of the overlap $ov = 3$.

Table 4

High-order finite-element discretizations for the 1D Poisson problem. Asymptotic convergence factors predicted by the two-grid analysis by using one smoothing step of the additive Schwarz method (AS) and the restricted additive Schwarz method (RAS) with natural weights. The size of the block is $k = p + 1$. Minimum overlap, $ov = 1$, is considered and the polynomial degree ranges from $p = 2$ to $p = 8$.

Smoother	Polynomial degree						
	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$
AS	0.50	0.09	0.17	0.11	0.17	0.12	0.16
RAS	0.50	0.12	0.15	0.11	0.17	0.12	0.16

Table 5

High-order finite-element discretizations for the 2D Poisson problem. Asymptotic convergence factors predicted by the two-grid analysis using two smoothing steps of the additive Schwarz method (AS) and the restricted additive Schwarz method (RAS) with natural weights. The size of the blocks is $k = (p + 1)^2$. Minimum overlap $ov = 1$ is assumed and the polynomial degree ranges from $p = 1$ to $p = 8$.

Smoother	Polynomial degree							
	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$
AS	0.14	0.15	0.21	0.25	0.31	0.36	0.40	0.43
RAS	0.19	0.15	0.21	0.25	0.31	0.36	0.40	0.43

view, as it avoids communication between processors. Next, we consider the performance of the multigrid method in terms of the number of smoothing steps at each level. In Table 6, we show the two-grid asymptotic convergence factors provided by LFA, ρ_{2g} , together with the asymptotic convergence factors computationally obtained by our own implementation, ρ_h , using $V(1,0)$, $V(1,1)$, $V(2,1)$ and $V(2,2)$ cycles. The coarsest grid used here within the V -cycle consists of only one inte-

Table 6

High-order finite-element discretizations for the 2D Poisson problem. Two-grid asymptotic convergence factors provided by LFA, ρ_{2g} , together with the computed asymptotic convergence factors, ρ_h , using $V(1,0)$, $V(1,1)$, $V(2,1)$, $V(2,2)$ cycles with the element-based restrictive additive Schwarz methods and natural weights. The polynomial degree ranges from $p = 1$ to $p = 8$.

p	$V(1,0)$		$V(1,1)$		$V(2,1)$		$V(2,2)$	
	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h
1	0.41	0.41	0.19	0.18	0.10	0.09	0.03	0.03
2	0.39	0.40	0.15	0.15	0.06	0.06	0.02	0.02
3	0.46	0.45	0.21	0.21	0.09	0.09	0.04	0.04
4	0.50	0.50	0.25	0.25	0.13	0.13	0.06	0.06
5	0.56	0.56	0.31	0.31	0.18	0.18	0.10	0.10
6	0.60	0.60	0.36	0.36	0.21	0.21	0.13	0.13
7	0.63	0.63	0.40	0.40	0.26	0.25	0.16	0.16
8	0.66	0.66	0.43	0.43	0.28	0.28	0.19	0.19

rior grid point. Again, we employ the element-based restrictive additive Schwarz smoother with natural weights for polynomial degree ranging from $p = 1$ to $p = 8$. We see that the asymptotic convergence factors predicted by the LFA match with high accuracy those computationally obtained by applying V -cycles.

5. Saddle point problems

In practice, Schwarz methods are among the most commonly used smoothers for solving saddle point problems. Whereas LFA has been widely applied to multiplicative Schwarz relaxations, few works on additive Schwarz smoothers are found in the literature. In [18], LFA was used to analyze this type of smoothers for the Stokes equations discretized by Taylor-Hood P2-P1 and Q2-Q1 schemes. This analysis can be seen as a particular case of the one introduced here, when using a

window of size 2×2 . In this section, we show that the presented Fourier analysis can be also applied to study the convergence of multigrid methods based on additive Schwarz methods applied to such saddle-point problems.

For this purpose, we consider the quasi-static Biot's model for poroelasticity. This model assumes that we have a deformable porous medium, which is linearly elastic, isotropic, homogeneous, and saturated by an incompressible Newtonian fluid. Given these assumptions, the well-known displacement-pressure formulation [27] must satisfy the following equations in a domain, $\Omega \subset \mathbb{R}^d$,

$$\begin{aligned} -\operatorname{div} \sigma' + \alpha \nabla p &= \rho \mathbf{g}, & \sigma' &= 2\mu \varepsilon(\mathbf{u}) + \lambda \operatorname{div}(\mathbf{u}), \\ \frac{\partial}{\partial t} \left(\frac{1}{M} p + \alpha \nabla \cdot \mathbf{u} \right) - \nabla \cdot \left(\frac{1}{\mu_f} \mathbf{K} (\nabla p - \rho_f \mathbf{g}) \right) &= f, \end{aligned} \quad (9)$$

where σ' and ε are the effective stress and strain tensors, λ and μ are the Lamé coefficients, p is the pore pressure, \mathbf{u} is the displacement vector field, \mathbf{g} is the gravity tensor, α is the Biot-Willis constant (which we assume is equal to one), \mathbf{K} is the permeability of the porous medium, μ_f is the fluid viscosity, M is the Biot modulus, and f is a source term. To complete the formulation of the problem, we add appropriate boundary and initial conditions. For instance,

$$\begin{aligned} p &= 0, & \sigma' \mathbf{n} &= \mathbf{t}, & \text{on } \Gamma_t, \\ \mathbf{u} &= \mathbf{0}, & \mathbf{K} (\nabla p - \rho_f \mathbf{g}) \cdot \mathbf{n} &= 0, & \text{on } \Gamma_c, \end{aligned} \quad (10)$$

where \mathbf{n} is the unit outward normal to the boundary and $\Gamma_t \cup \Gamma_c = \Gamma := \partial\Omega$, with Γ_t and Γ_c disjoint subsets of Γ having non null measure. For the initial time, $t = 0$, the following condition is fulfilled,

$$\left(\frac{1}{M} p + \alpha \nabla \cdot \mathbf{u} \right) (\mathbf{x}, 0) = 0, \quad \mathbf{x} \in \Omega. \quad (11)$$

To discretize the problem, we use Taylor-Hood finite elements in space, $\mathbb{Q}_2 - \mathbb{Q}_1$, and a backward-Euler scheme in time. Let \mathcal{T}_h be a triangulation of Ω composed of rectangles. Defining the discrete spaces as

$$\mathbf{V}_h = \{ \mathbf{u}_h \in (H^1(\Omega))^d \mid \forall T \in \mathcal{T}_h, \mathbf{u}_h|_T \in \mathbb{Q}_2^d, \mathbf{u}_h|_{\Gamma_c} = \mathbf{0} \},$$

$$\mathcal{Q}_h = \{ p_h \in H^1(\Omega) \mid \forall T \in \mathcal{T}_h, p_h|_T \in \mathbb{Q}_1, p_h|_{\Gamma_t} = 0 \},$$

the fully discretized scheme at time t_m , $m = 1, 2, \dots$, is written as follows: find $(\mathbf{u}_h^m, p_h^m) \in \mathbf{V}_h \times \mathcal{Q}_h$ such that

$$a(\mathbf{u}_h^m, \mathbf{v}_h) - \alpha(p_h^m, \operatorname{div} \mathbf{v}_h) = (\rho \mathbf{g}, \mathbf{v}_h), \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad (12)$$

$$\alpha(\operatorname{div} \bar{\partial}_t \mathbf{u}_h^m, q_h) + b(p_h^m, q_h) = (f_h^m, q_h) + (\mathbf{K} \mu_f^{-1} \rho_f \mathbf{g}, \nabla q_h), \quad \forall q_h \in \mathcal{Q}_h, \quad (13)$$

where $\bar{\partial}_t \mathbf{u}_h^m := (\mathbf{u}_h^m - \mathbf{u}_h^{m-1})/\tau$, with τ the time discretization parameter, (\cdot, \cdot) is the standard inner product in the space $L_2(\Omega)$, and the bilinear forms $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ are given as

$$a(\mathbf{u}, \mathbf{v}) = 2\mu \int_{\Omega} \varepsilon(\mathbf{u}) : \varepsilon(\mathbf{v}) d\Omega + \lambda \int_{\Omega} \operatorname{div} \mathbf{u} \operatorname{div} \mathbf{v} d\Omega,$$

$$b(p, q) = \int_{\Omega} \frac{\mathbf{K}}{\mu_f} \nabla p \cdot \nabla q d\Omega.$$

This fully discrete scheme leads to a saddle-point system of equations at each time step of the form $\mathcal{A}x = b$, where matrix \mathcal{A} is a 2×2 block symmetric indefinite matrix:

$$\mathcal{A} = \begin{pmatrix} A & B^T \\ B & -C \end{pmatrix}, \quad (14)$$

with matrices A and C both being symmetric and positive definite.

To solve this system of equations, we consider a geometric multigrid method with standard coarsening based on an additive Schwarz method as the relaxation scheme. For transfer operators we consider the canonical restriction and prolongation operators. The classical Schwarz smoother for this type of saddle-point problems is typically defined as a




Fig. 2. Minimum size window for the two-dimensional Biot's model. The block centered on the upper-left pressure unknown of the grid is shown in blue.

Table 7

Taylor-Hood discretizations of the 2D quasi-static Biot's model. Two-grid asymptotic convergence factors provided by LFA, ρ_{2g} , together with the computed asymptotic convergence factors, ρ_h , using several permeability values and different numbers of smoothing steps of the 51-point additive Schwarz relaxation with natural weights.

K	$W(1,0)$		$W(1,1)$		$W(2,1)$		$W(2,2)$	
	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h	ρ_{2g}	ρ_h
1	0.49	0.49	0.25	0.22	0.12	0.11	0.06	0.06
10^{-3}	0.49	0.49	0.25	0.21	0.12	0.11	0.06	0.06
10^{-6}	0.49	0.49	0.25	0.21	0.12	0.11	0.06	0.06
10^{-9}	0.65	0.65	0.42	0.42	0.27	0.27	0.18	0.18
10^{-12}	0.72	0.72	0.52	0.52	0.52	0.56	0.28	0.28
10^{-15}	0.72	0.72	0.52	0.52	0.52	0.56	0.28	0.28

set of blocks that consists of one pressure unknown and all the velocity unknowns that are connected to it. That is, the degrees of freedom corresponding to the nonzero entries in the i th row of B plus the i -th pressure degree of freedom. This yields a local system of size 51×51 and 76×76 for two-dimensional and three-dimensional problems, respectively. Due to the high computational cost associated with this type of smoother, the parallelization of the relaxation method becomes crucial for real applications. Therefore, the additive and the restricted additive Schwarz smoothers are a more natural choice than multiplicative versions, and it is clear that a Fourier analysis tool for them is of great importance for the design of geometric multigrid methods for saddle point problems.

For two-dimensional problems, since the number of different stencils is two, and each basis function for the pressure shares common support with five displacement unknowns on each direction, the minimum window size required for the analysis is 8×8 , so we choose that size here. In Fig. 2, we show the periodic extension of this window with a selected block of unknowns in blue corresponding to the upper-left pressure unknown on the computational domain.

In the following numerical results, we fix the Lamé parameters in such a way that the Young's modulus is $E = 3 \times 10^4$, and the Poisson's ratio is $\nu = 0.2$. We consider different values for permeability, K , ranging from $K = 1$ to $K = 10^{-15}$. First, we consider the multigrid algorithm based on additive Schwarz smoothers using the natural weights, i.e., $w = 1$ for the pressure and $w = 1/9$, $1/6$, or $w = 1/4$ for the displacement depending on if the degree of freedom corresponds to a vertex, to an edge, or a cell node, respectively. In Table 7, we show the asymptotic convergence factors provided by our analysis together with the numerical ones provided by our implementation using W-cycles. We consider different numbers of smoothing steps and different values of permeability.

In Table 7, we observe that there is a good match between the factors provided by LFA and the ones experimentally obtained. We also see that the performance of the multigrid method deteriorates when K

Table 8

Taylor-Hood discretizations of the 2D quasi-static Biot's model. Two-grid asymptotic convergence factors provided by LFA, using several permeability values and different numbers of smoothing steps of the 51-point additive Schwarz relaxation with weights (0.09, 0.22, 1.02).

K	$W(1,0)$	$W(1,1)$	$W(2,1)$	$W(2,2)$
1	0.58	0.34	0.19	0.11
10^{-3}	0.58	0.34	0.19	0.11
10^{-6}	0.58	0.34	0.19	0.11
10^{-9}	0.58	0.34	0.19	0.11
10^{-12}	0.60	0.36	0.21	0.13
10^{-15}	0.60	0.36	0.21	0.13

Table 9

Taylor-Hood discretizations of the 2D quasi-static Biot's model. Number of multigrid iterations using the additive Schwarz smoothers required to reduce the initial residual by a factor of $tol = 10^{-10}$, using the weights (0.09, 0.22, 1.02) with V - and W -cycles for several smoothing steps and permeabilities.

K	Smoothing steps (v_1, v_2)							
	(1,0)		(1,1)		(2,1)		(2,2)	
	V	W	V	W	V	W	V	W
1	39	40	20	20	13	14	10	10
10^{-3}	40	40	20	20	14	13	10	10
10^{-6}	39	40	20	20	13	14	10	10
10^{-9}	40	40	23	20	14	13	10	10
10^{-12}	50	43	40	22	18	15	12	12
10^{-15}	50	45	40	23	18	15	12	12

tends to zero, yielding a nonrobust algorithm. However, an additional advantage of our analysis is that we are able to find the optimal weights for a fixed permeability in order to obtain a robust algorithm. Since permeability is usually heterogeneous in real applications, we choose the optimal weights corresponding to the worst case scenario ($K = 0$), that is, $w = 0.09$ for the displacements at vertices and edges, $w = 0.22$ for the displacements at interior points, and $w = 1.02$ for the pressure variables. In Table 8, we show the asymptotic convergence factors applying the additive Schwarz smoother with these optimal weights provided by our analysis, considering different number of smoothing steps and several permeability values.

Given that V -cycles are less expensive than W -cycles, we would like to compare their performance in number of iterations. Hence, in Table 9, we show the number of iterations, n_{iter} , used in our implementation required to reduce the initial residual by a factor of $tol = 10^{-10}$, using the optimal weights with V - and W -cycles for several smoothing steps and permeabilities.

We conclude that V -cycles are also robust with respect to K when applying $V(2,1)$ - or $V(2,2)$ -cycles since both cycles yield a similar number of iterations in order to reach the stopping criterion. Thus, the use of $V(2,2)$ -cycles with the 51-point additive Schwarz smoother applying the weights (0.09, 0.22, 1.02) is a good approach for solving the two-field formulation of Biot's equations.

6. Conclusions

A general framework for the analysis of multigrid methods based on additive Schwarz relaxations is presented in this work. This approach is based on a local Fourier analysis, which considers a basis of the Fourier space different from the classical one given in terms of the Fourier modes. The proposed analysis allows the study of a class of additive Schwarz smoothers for any problem and discretization. As examples, high-order finite-element discretizations of Poisson's problem and saddle point problems are considered since these are two formulations

that benefit from applying these relaxation procedures. In addition, both additive and restricted additive Schwarz smoothers are considered. In all cases, we show an excellent match between the convergence rates provided by the analysis and those asymptotic convergence factors computationally obtained from a multigrid implementation. In the case of high-order finite-element discretizations of the Poisson problem, for each polynomial degree p , we find that the restricted additive Schwarz smoother achieves comparable convergence rates to the additive Schwarz smoother, having better favorable properties in terms of scalability and applicability to high-performance computing. Thus, we conclude that the restricted additive Schwarz method can be a good alternative as a relaxation procedure in a multigrid method for solving high-order discretizations of elliptic equations. Regarding the application to saddle-point problems, we consider Biot's model for poroelasticity as a model problem and show that the proposed analysis allows one to obtain optimal weights to define a multigrid solver based on additive Schwarz smoothers, which is robust with respect to physical parameters involved in the model. The proposed analysis, however, is a general approach that can be applied to any type of discretization and problem.

Data availability

No data was used for the research described in the article.

Acknowledgements

The work of Álvaro Pé and Carmen Rodrigo is supported in part by the Spanish Grant PID2022-140108NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by "ERDF A way of making Europe", and by the "European Union NextGenerationEU/PRTR". Francisco J. Gaspar has received funding from the Spanish Grant PID2019-105574GB-I00 funded by MCIN/AEI/10.13039/501100011033 and by "ERDF A way of making Europe", and by the "European Union NextGenerationEU/PRTR". Álvaro Pé, Carmen Rodrigo and Francisco J. Gaspar acknowledge support from the Diputación General de Aragón, Spain (Grupo de referencia APEDIF, ref. E24_17R). The work of James Adler and Xiaozhe Hu is partially supported by the National Science Foundation (NSF) under Grant DMS-2208267. The research of Ludmil Zikatanov is supported in part by the U.S.-Norway Fulbright Foundation and the U.S. National Science Foundation Grant DMS-2208249.

References

- [1] S. Vanka, Block-implicit multigrid solution of Navier-Stokes equations in primitive variables, *J. Comput. Phys.* 65 (1) (1986) 138–158, [https://doi.org/10.1016/0021-9991\(86\)90008-2](https://doi.org/10.1016/0021-9991(86)90008-2), <https://www.sciencedirect.com/science/article/pii/0021999186900082>.
- [2] J. Schöberl, W. Zulehner, On Schwarz-type smoothers for saddle point problems, *Numer. Math.* 95 (2) (2003) 377–399, <https://doi.org/10.1007/s00211-002-0448-3>.
- [3] X.-C. Cai, M. Sarkis, A restricted additive Schwarz preconditioner for general sparse linear systems, *SIAM J. Sci. Comput.* 21 (2) (1999) 792–797, <https://doi.org/10.1137/S106482759732678X>.
- [4] A. Frommer, D.B. Szyld, An algebraic convergence theory for restricted additive Schwarz methods using weighted max norms, *SIAM J. Numer. Anal.* 39 (2) (2001) 463–479, <https://doi.org/10.1137/S0036142900370824>.
- [5] S. Saberi, G. Meschke, A. Vogel, A restricted additive Vanka smoother for geometric multigrid, *J. Comput. Phys.* 459 (2022) 111123, <https://doi.org/10.1016/j.jcp.2022.111123>, <https://www.sciencedirect.com/science/article/pii/S0021999122001851>.
- [6] A. Brandt, Multi-level adaptive solutions to boundary-value problems, *Math. Comput.* 31 (138) (1977) 333–390.
- [7] A. Brandt, Rigorous quantitative analysis of multigrid, I: constant coefficients two-level cycle with L2-norm, *SIAM J. Numer. Anal.* 31 (6) (1994) 1695–1730.
- [8] K. Stüben, U. Trottenberg, Multigrid methods: fundamental algorithms, model problem analysis and applications, in: W. Hackbusch, U. Trottenberg (Eds.), *Multigrid Methods*, Springer Berlin Heidelberg, Berlin, Heidelberg, 1982, pp. 1–176.
- [9] P. Wesseling, *An Introduction to Multigrid Methods*, R.T. Edwards, 2004, <https://books.google.es/books?id=RoRGAAAYAAJ>.
- [10] U. Trottenberg, C. Oosterlee, A. Schüller, *Multigrid*, Elsevier Science, 2001, https://books.google.es/books?id=-og1wD-Nx_wC.

- [11] R. Wienands, W. Joppich, *Practical Fourier Analysis for Multigrid Methods*, Chapman and Hall/CRC Press, 2005.
- [12] C. Rodrigo, F. Gaspar, L. Zikatanov, On the validity of the local Fourier analysis, *J. Comput. Math.* 37 (3) (2018) 340–348, <https://doi.org/10.4208/jcm.1803-m2017-0294>, http://global-sci.org/intro/article_detail/jcm/12725.html.
- [13] S. Sivaloganathan, The use of local mode analysis in the design and comparison of multigrid methods, *Comput. Phys. Commun.* 65 (1) (1991) 246–252, [https://doi.org/10.1016/0010-4655\(91\)90178-N](https://doi.org/10.1016/0010-4655(91)90178-N), <https://www.sciencedirect.com/science/article/pii/001046559190178N>.
- [14] J. Molenaar, *A Two-Grid Analysis of the Combination of Mixed Finite Elements and Vanka-Type Relaxation*, Birkhäuser, Basel, Basel, 1991, pp. 313–323.
- [15] C. Rodrigo, F. Gaspar, F. Lisbona, On a local Fourier analysis for overlapping block smoothers on triangular grids, *Appl. Numer. Math.* 105 (2016) 96–111, <https://doi.org/10.1016/j.apnum.2016.02.006>, <https://www.sciencedirect.com/science/article/pii/S0168927416300022>.
- [16] S.P. MacLachlan, C.W. Oosterlee, Local Fourier analysis for multigrid with overlapping smoothers applied to systems of PDEs, *Numer. Linear Algebra Appl.* 18 (4) (2011) 751–774, <https://doi.org/10.1002/nla.762>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/nla.762>.
- [17] T. Boonen, J.V. Lent, S. Vandewalle, Local Fourier analysis of multigrid for the curl-curl equation, *SIAM J. Sci. Comput.* 30 (2008) 1730–1755.
- [18] P.E. Farrell, Y. He, S.P. MacLachlan, A local Fourier analysis of additive Vanka relaxation for the Stokes equations, *Numer. Linear Algebra Appl.* 28 (3) (2021) e2306, <https://doi.org/10.1002/nla.2306>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/nla.2306>.
- [19] C. Greif, Y. He, A closed-form multigrid smoothing factor for an additive Vanka-type smoother applied to the Poisson equation, *Numer. Linear Algebra Appl.* 30 (2023) e2500, <https://doi.org/10.1002/nla.2500>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/nla.2500>.
- [20] P. Kumar, C. Rodrigo, F.J. Gaspar, C.W. Oosterlee, On local Fourier analysis of multigrid methods for PDEs with jumping and random coefficients, *SIAM J. Sci. Comput.* 41 (3) (2019) A1385–A1413, <https://doi.org/10.1137/18M1173769>.
- [21] J. Brown, Y. He, S. MacLachlan, Local Fourier analysis of balancing domain decomposition by constraints algorithms, *SIAM J. Sci. Comput.* 41 (5) (2019) S346–S369, <https://doi.org/10.1137/18M1191373>.
- [22] M. Bolten, H. Rittich, Fourier analysis of periodic stencils in multigrid methods, *SIAM J. Sci. Comput.* 40 (3) (2018) A1642–A1668, <https://doi.org/10.1137/16M1073959>.
- [23] D. Greenfeld, M. Galun, R. Basri, I. Yavneh, R. Kimmel, Learning to optimize multigrid PDE solvers, in: K. Chaudhuri, R. Salakhutdinov (Eds.), *Proceedings of the 36th International Conference on Machine Learning*, in: *Proceedings of Machine Learning Research*, PMLR, vol. 97, 2019, pp. 2415–2423.
- [24] C. Rodrigo, F. Gaspar, F. Lisbona, Multicolor Fourier analysis of the multigrid method for quadratic FEM discretizations, *Appl. Math. Comput.* 218 (22) (2012) 11182–11195, <https://doi.org/10.1016/j.amc.2012.05.008>, <https://www.sciencedirect.com/science/article/pii/S0096300312004936>.
- [25] Álvaro Pé de la Riva, C. Rodrigo, F.J. Gaspar, A two-level method for isogeometric discretizations based on multiplicative Schwarz iterations, *Comput. Math. Appl.* 100 (2021) 41–50, <https://doi.org/10.1016/j.camwa.2021.08.020>, <https://www.sciencedirect.com/science/article/pii/S0898122121003126>.
- [26] F. Gaspar, Code for manuscript entitled “A local Fourier analysis for additive Schwarz smoothers”, <https://doi.org/10.5281/zenodo.10405571>, Dec. 2023.
- [27] H.F. Wang, *Theory of Linear Poroelasticity with Applications to Geomechanics and Hydrogeology*, Princeton University Press, 2000, <http://www.jstor.org/stable/j.ctt1jktrr4>.